

Lizard fauna (Squamata, Sauria) from Serra do Ouro Branco, southern Espinhaço Range, Minas Gerais, Brazil

Antônio Jorge do Rosário Cruz^{1,2*}, Leandro de Oliveira Drummond^{2,3}, Virginia Duarte Lucena², Adriele Prisca de Magalhães^{1,2}, Caryne Aparecida de Carvalho Braga^{1,2,3}, Jaqueline Malta Rolin² and Maria Rita Silvério Pires^{1,2}

1 Universidade Federal de Ouro Preto, Campus Morro do Cruzeiro, Programa de Pós-graduação em Ecologia de Biomas Tropicais. CEP 35400-000. Ouro Preto, MG, Brasil.

2 Universidade Federal de Ouro Preto, Campus Morro do Cruzeiro, Departamento de Evolução, Biodiversidade e Meio Ambiente. CEP 35400-000. Ouro Preto, MG, Brasil.

3 Universidade Federal do Rio de Janeiro, Departamento de Ecologia, Laboratório de Vertebrados. CP 68020, Ilha do Fundão. CEP 21941-901, Rio de Janeiro, RJ, Brazil.

* Corresponding author. E-mail: cruzony@gmail.com

ABSTRACT: The present study evaluated the lizard fauna in Serra do Ouro Branco, Minas Gerais, Brazil, a transition area between the Atlantic Forest and Cerrado. Data was collected using pitfall traps, active and occasional encounters, and through information from zoological collections and the literature. Field sampling was performed in two stages over a period of 36 months: from December 2006 to December 2008, and from January to December 2010. The study area is home to 15 species belonging to eight families: Anguidae, Gekkonidae, Gymnophthalmidae, Leiosauridae, Polychrotidae, Mabuyidae, Teiidae, and Tropiduridae. Lizard species evaluated in this study represent a relevant asset for this zoological group inhabiting the transitional areas between the Atlantic Forest and Cerrado. They include strictly forest species and endemic Atlantic Forest species such as *Ecpleopus gaudichaudii*, *Enyalius perditus* and *Heterodactylus imbricatus* in addition to typical Cerrado taxa common to both biomes.

DOI: 10.15560/10.6.1290

INTRODUCTION

Lizards are traditionally studied as a cohesive group, although recognized as paraphyletic (Pyron *et al.* 2013). These organisms are considered model organisms, either in studies of human impact on wildlife (Kitchener *et al.* 1980; Gibbons *et al.* 2000; Lehtinen *et al.* 2003; Jellinek *et al.* 2004; Bell and Donnelly 2006; Gardner *et al.* 2007; Dixo and Martins 2008; Dixo and Metzger 2009), or in the interpretation of data on patterns of distribution and endemism between ecosystems (Pianka 1973; Vanzolini *et al.* 1980; Huey *et al.* 1983; Rodrigues 1987, 2005; Recorder and Nogueira 2007). They comprise one of the most diverse groups of amniotes, with about 5800 described species (Uetz and Hosek 2013), of which 248 occur in Brazil (Bérnils and Costa 2012). However, this great diversity is threatened in several ecosystems, mainly due habitat loss (Gibbons *et al.* 2000). In this aspect, the Brazilian state of Minas Gerais deserves special attention, since it contains two biodiversity hotspots (Cerrado and Atlantic Rain Forest; see Mittermeier *et al.* 2004).

About one quarter of all Brazilian lizards were reported for the state of Minas Gerais (Bérnils *et al.* 2009). However, basic information, such as area of occurrence and studies on ecology and natural history, is still lacking for most species. Local composition of lizard communities has been surveyed only in the following localities: Lagoa Santa (Reinhardt and Lutken 1861., based on specimens from Lagoa Santa collected by Peter Lund); Northeast of Minas Gerais (Feio and Caramaschi 2002); Parque Nacional Grande Sertão Veredas (Recorder and Nogueira 2007);

Estação de Preservação e Desenvolvimento Ambiental de Peti (Bertoluci *et al.* 2009); Viçosa (Costa *et al.* 2009); Ritópolis (Sousa *et al.* 2010); Juiz de Fora (Sousa *et al.* 2012); Serra do Brigadeiro (Moura *et al.* 2012); Reserva Ecológica Unilavras-Boqueirão (Novelli *et al.* 2012); and Instituto Inhotim (Linares and Eterovick 2013).

The Espinhaço Range is a mountainous formation that extends about 1200 km from the state of Minas Gerais to the state of Bahia, containing ecosystems such as Rocky Grasslands (Campos Rupestres), Cerrado, Atlantic Rainforest and Caatinga (Derby 1906; Giulietti *et al.* 1997; Vitta 2002; Almeida-Abreu and Renger 2002). Due to the high degree of endemism of the fauna and flora, and human disturbance, this mountain range, including the Serra do Ouro Branco, was declared a priority area for Biodiversity conservation in Minas Gerais (Drummond *et al.* 2005).

Most of the biodiversity of the Espinhaço Range is yet unsampled, besides its great biological relevance. The reptile fauna is particularly poorly known. Most reptile studies in the southern parts of the range have examined only the snake communities in the Serra do Cipó (Assis *et al.* 1999); Serra do Ouro Branco (São Pedro and Pires 2009) and the municipalities of Ouro Preto, Itabirito and Mariana (Silveira *et al.* 2010). Surveys of lizard species were conducted only in Chapada Diamantina by Juncá (2005), and in EPDA Peti by Bertoluci *et al.* (2009).

Recently the Parque Estadual Serra do Ouro Branco was created to protect the biodiversity of this mountainous complex in the southern end of the Espinhaço Range, state of Minas Gerais. However there is little information




FIGURE 1: Location of the Serra do Ouro Branco, state of Minas Gerais, Brazil.

about the composition, richness or distribution of the lizard fauna of this region, making it difficult to evaluate the importance of this protected area for the conservation of this group. Thus, the present study aims to provide an inventory and a key for identification of the lizards from the Serra de Ouro Branco.

MATERIALS AND METHODS

Study Area

This study was carried out in the Serra do Ouro Branco ($20^{\circ} 31' 15''$ S, $43^{\circ} 41' 31''$ W), located in the municipality of Ouro Branco, state of Minas Gerais, in the southern end of the Espinhaço Range (Figure 1), including the area surrounding Itatiaia, a village with 18th century buildings, today with about 80 residences. The elevation of the region varies between 900 m to 1600 m (Veloso *et al.* 1991). The vegetation of the region consists of Seasonal Semideciduous Forest and Rocky Grassland at higher elevations, plus grasslands and other environments typical of the Cerrado biome (Veloso *et al.* 1991). The climate is Cwb mesothermal, according to Köppen's classification, with a dry season between the months of June and September (winter), and a higher incidence of rainfall from November to February. The mean annual temperature is 20.7°C and the mean annual rainfall is 1321.07 mm, according to data from the Meteorological Station of Gerdaú Açominas, located approximately 15 km from the study area.

Sampling

Fieldwork was carried out in two stages, totaling 36 months of sampling. To better cover the different environments of Serra do Ouro Branco, three sampling

methods were used: pitfall traps (Cechin and Martins 2000), active search (Crump and Scott Jr. 1994) and incidental encounters. Animals were collected under permits issued by the Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis- IBAMA-481/06-NUFAS-MG and Instituto Chico Mendes de Conservação da Biodiversidade- ICMBIO- 21543-1.

During the first stage, from December 2006 to December 2008, pitfall traps were installed in five distinct forest areas. Each row consisted of ten 60 L buckets, buried at intervals of four m in a straight line. The buckets were connected by one m high fences. Every month, the traps remained open for five consecutive days, giving a total sampling effort of 50 buckets per day for a total of 125 sampling days, or 6250 bucket-nights.

During the second stage, from January to December 2010, a new sample design was adopted, in which four 60 L buckets were buried in a Y-shaped array *i.e.*, one central bucket linked to three peripheral ones by 1 m high fences, the three arms forming angles of approximately 120° . Thus, 27 new sampling points were installed in sets of three, in nine different areas. The traps remained open for five days/month, giving a total sampling effort of 108 buckets per day for a total of 60 days of sampling, or 6480 bucket-nights. This type of trap was used exclusively in forest environments, where it was possible to find soil deep enough to bury the buckets.

The active search method (Crump and Scott Jr. 1994) was used in grasslands, rocky grasslands, gallery forests and forests. This method was carried out in nonconsecutive diurnal (09:00–15:00 h) and nocturnal (18:00–22:00 h) surveys, by two or more researchers. Specimens found while walking along the trails between the pitfall traps

were considered incidental records.

The specimens were killed using 0.5% lidocaine, fixed in 10% formalin, preserved in 70% ethanol, and deposited in the herpetological collection of the Laboratório de Zoologia dos Vertebrados of the Universidade Federal de Ouro Preto (LZV-UFOP). Field guides, identification keys and articles were used to assist in species identification (Vanzolini 1948; Peters and Donoso-Barros 1970; Vanzolini 1978; Vanzolini *et al.* 1980; Rodrigues 1987; Avila-Pires 1995; Doan 2003; Rodrigues *et al.* 2007, 2009). Taxonomy follows Estes *et al.* (1988), Frost *et al.* (2001) and Gamble *et al.* (2008). For the Teiidae we follow the recommendations of Pyron *et al.* (2013) to maintain the traditional nomenclature and for the Scincidae we follow the nomenclature proposed by Hedges and Conn (2012).

To supplement the samples, specimens from the herpetological collection of the LZV-UFOP were examined as well as one specimen from the herpetological collection of the Museu de Zoologia João Moojen of the Universidade Federal de Viçosa (MZUFV), Minas Gerais.

To evaluate the effectiveness of the sampling to detect the lizard richness of the Serra do Ouro Branco, we constructed an individual based rarefaction curve (Colwell 2011), using the program EstimateS 9.1 (Colwell 2013). We contextualize regionally the local lizard composition of the Serra do Ouro Branco evaluating the dissimilarity

between it and other sampled areas in Minas Gerais. The dissimilarity of lizard communities of Minas Gerais was evaluated using Jaccard’s distance coefficient, a method based only on the composition, which does not take into account the species abundance. The Jaccard’s coefficient was used to construct a dendrogram based on the UPGMA grouping method (Unweighted Pair Group Method with Arithmetic Mean). This analysis was performed using the vegan package (Oksanen *et al.* 2010) at R software (R Development Core Team 2013). For the analyses, we exclude species identified only to the genus taxonomic level.

RESULTS

A total of 15 lizard species were recorded in the Serra do Ouro Branco, belonging to 13 genera and eight families. Gymnophthalmidae and Leiosauridae, represented by three species each, were the most diverse families in the area, followed by Mabuyidae, Teiidae and Tropiduridae, represented by two species each, and Anguidae, Gekkonidae and Polychrotidae represented by a single species each. Detailed accounts of habits, habitats and distribution of each species are in Table 1. Photographs of all species are in Figures 2 and 3. Voucher specimens are listed in Appendix 1, and scientific articles, books and guides consulted to analyze the occurrence of the lizards in

TABLE 1: List of species of lizards of the Serra do Ouro Branco, state of Minas Gerais, recorded between December 2006 and December 2008 and from January to December 2010. **Activity:** D, diurnal; N, nocturnal. **Habit:** SF, semi-fossorial; TR, terrestrial; SA, semi-arboreal; A, arboreal; C, climber; FL, forests; CL, clearings; RG, rocky grasslands (*Campos Rupestres*); UA, urban areas. **Biomes:** AR, Atlantic Rainforest; CE, Cerrado; WD, wide distribution (species occurring in more than two biomes). **Methodology** (Capture): PT, pitfall; AS, active search; OC, incidental encounter.

TAXON	ACTIVITY	HABIT	HABITAT				BIOMES	METHODOLOGY
			FL	CL	CR	UA		
Anguidae								
<i>Ophiodes striatus</i> (SPIX, 1824)	D	SF	x	x	—	x	WD	PT/OC
Gekkonidae								
<i>Hemidactylus mabouia</i> (Moreau de Jonnès, 1818)	N	C	—	—	—	x	WD	OC
Gymnophthalmidae								
<i>Cercosaura quadrilineatus</i> (Boettger, 1876)	D	SF	x	—	—	—	AR	PT
<i>Ecpleopus gaudichadii</i> Duméril & Bibron, 1839	D	SF	x	—	—	—	AR	PT
<i>Heterodactylus imbricatus</i> Spix, 1825	D	SF	x	—	—	—	AR	PT
Leiossauridae								
<i>Enyalius perditus</i> Jackson, 1978	D	SA	x	—	—	—	AR	PT/AS/OC
<i>Enyalius bilineatus</i> Duméril & Bibron, 1837	D	SA	x	—	—	—	AR/CE	PT/OC
<i>Urostrophus vaultieri</i> Duméril & Bibron, 1837	D	SA	x	—	—	—	AR	PT
Polychrotidae								
<i>Polychrus acutirostris</i> Spix, 1825	D	A	x	—	—	—	WD	OC
Mabuyidae								
<i>Aspronema dorsivittatum</i> (Cope,1862)	D	T	—	—	x	—	WD	OC
<i>Notomabuya frenata</i> (Cope, 1862)	D	T	x	—	—	x	WD	PT/OC
Teiidae								
<i>Ameiva ameiva</i> (Linnaeus , 1758)	D	T	—	—	x	—	WD	PT/AS/OC
<i>Tupinambis merianae</i> (Duméril & Bibron, 1839)	D	T	x	x	x	—	WD	PT/OC
Tropiduridae								
<i>Tropidurus itambere</i> Rodrigues, 1987	D	T	—	—	x	x	CE	AS
<i>Tropidurus torquatus</i> Wiegmann, 1934	D	T	—	x	—	x	AR/CE	AS/OC




FIGURE 2: Lizards from Serra do Ouro Branco: A) *Ophiodes striatus*; B) *Hemidactylus mabouia* (Photo: Pedro H. Bernardo); C) *Cercosaura quadrilineatus*; D) *Ecpleopus gaudichaudii*; E) *Heterodactylus imbricatus*; F) *Enyalius perditus* (male); G) *Enyalius perditus* (Juvenile); H) *Enyalius bilineatus*.




FIGURE 3: Lizards from Serra do Ouro Branco: A) *Urostrophus vautieri*; B) *Polychrus acutirostris* (Photo: Pedro H. Bernardo); C) *Notomabuya frenata*; D) *Aspronema dorsivittatum* (Photo: Henrique C. Costa); E) *Ameiva ameiva* (Photo: Pedro H. Bernardo); F) *Tupinambis merianae*; G) *Tropidurus torquatus* (photo: Pedro Bernardo); H) *Tropidurus itambere*.

different biomes are listed in Appendix 2. The rarefaction curve based on the number of individuals recorded from all sampling methods has not stabilized, indicating the possibility presence of species not yet recorded in the study area (Figure 4).

The lizard fauna of Serra do Ouro Branco is most similar to those communities of the Atlantic Rainforest in Juiz de Fora and the Serra do Brigadeiro (dissimilarity coefficients of 0.45 and 0.47 respectively). In fact, the lizard fauna of Serra do Brigadeiro is a subset of the species founded in Serra de Ouro Branco. The lizard faunas of the farthest areas, Northeast of Minas Gerais and Parque Nacional Grande Sertão Veredas, showed the lowest similarity with that of the present study (dissimilarity coefficients of 0,89 and 0,78, respectively). The resulting dendrogram of dissimilarity reinforces the greater similarity of lizard fauna of the Serra do Ouro Branco with those of other areas in the Atlantic Rainforest of southeastern Minas Gerais (Juiz de Fora, Serra do Brigadeiro, Viçosa and EPDA-Peti; Figure 5).

DISCUSSION

The number of lizard species recorded for Serra do Ouro Branco corresponds to 26.31% of the lizards known for the state of Minas Gerais (Bérnils *et al.* 2009) and 6% of the species of Brazilian lizards (Bérnils and Costa 2012). Colli *et al.* (2002) and Nogueira (2006) state that in well-sampled localities of the Cerrado the estimated richness of lizards is 13 to 28 species. There are no studies providing such estimates for the Atlantic Rainforest. However, some studies carried out specifically in this biome found lizard communities ranging from four to 15 species (*e.g.*, Bertoluci *et al.* 2009; Carvalho *et al.* 2007; Cicchi *et al.* 2009; Condez *et al.* 2009; Costa *et al.* 2009; Dixo and Verdade 2006; Forlani *et al.* 2010; Moura *et al.* 2012). Thus Serra de Ouro Branco has both lower diversity than other studies in the Cerrado and higher diversity than studies in Atlantic Rainforest. Nevertheless, these comparisons are not conclusive, since differences in the extent of the sampling areas and the methodologies used in each study




FIGURE 4: Individual sample based rarefaction curve of the species recorded in Serra do Ouro Branco, Minas Gerais.

site can directly influence species richness (Rodrigues 2005).

The use of different methods of capture proved to be essential to obtain a more complete picture of the assemblage of lizards of Serra do Ouro Branco. Pitfall traps were responsible for the capture of the majority of the species (10 species, including 5 species exclusively. However, active search and incidental encounters together added five species to the list: *Hemidactylus mabouia*, *Polychrus acutirostris*, *Aspronema dorsivittatum*, *Tropidurus torquatus*, and *Tropidurus itambere*. These species are good climbers, able to escape from the buckets, and are typical of open environments, which cannot be sampled by pitfalls since the soil is shallow.

All native species collected are diurnal. The only nocturnal lizard present in the region is the gekkonid *Hemidactylus mabouia*. Individuals of this species were observed near light sources in human settlements, possibly preying on small invertebrates. This gecko is an exotic (African) species and a successful colonizer with a broad distribution in the tropics (Rödder *et al.* 2008).




FIGURE 5: Dissimilarity dendrogram of the composition of lizard species in 11 localities from the state of Minas Gerais, Brazil. UPGMA cluster method and Jaccard's dissimilarity index. 1- Serra do Ouro Branco (this study), 2- EPDA Peti (Bertoluci *et al.* 2009), 3- Viçosa (Costa *et al.* 2009), 4- Serra do Brigadeiro (Moura *et al.* 2012), 5- Juiz de Fora (Sousa *et al.* 2012), 6- Ritópolis (Sousa *et al.* 2010), 7- Parque Nacional Grande Sertão Veredas (Recorder and Nogueira 2007), 8- Reserva Ecológica Unilavras-Boqueirão (Novelli *et al.* 2012), 9- Lagoa Santa (Reinhardt and Lutken 1861), 10- Nordeste de Minas Gerais (Feio and Caramaschi 2002), 11- Instituto Inhotim (Linares and Eterovick 2013).

Although it may be abundant in natural environments of various biomes, this species is primarily associated with urban areas (Rocha et al. 2011). In Serra do Ouro Branco *H. mabouia* is apparently restricted to perianthropic areas. Due to the differential daily activity and the specificity to anthropogenic habitats is unlikely that this species is competing directly with native species.

The majority of the lizard fauna of Serra do Ouro Branco is composed of species with wide distributions. About 60% (nine species) of the lizards recorded in the Serra do Ouro Branco have distributions covering both Atlantic Rainforest and Cerrado habitats, and seven of these species have distribution covering more than two biomes. However, some of these species are known to be species complexes, such as *Ophiodes striatus* (Borges-Martins 1998) and *Enyalius bilineatus* (see Rodrigues et al. 2006), and the populations of the Serra de Ouro Branco may actually belong to species with more restricted distribution.

The regional lizard fauna is composed predominantly of species with wide distribution and endemic species from the Atlantic Rainforest biomes. Despite the existence of records of *Heterodactylus imbricatus* (Novelli et al. 2011) and *Urostrophus vautieri* (Novelli et al. 2012) in Cerrado, we still consider these species as endemic to the Atlantic Forest biome, since these records are from riparian forests in marginal areas of this biome. Considering this, there are five species endemic from Atlantic Rainforest in the Serra do Ouro Branco. Added to this, Jaccard's dissimilarity index reinforces the greater influence of the Atlantic Rainforest in the composition of the regional fauna, grouping the Serra de Ouro Branco to areas of this biome in southeastern Minas Gerais. However, this apparent low influence of the Cerrado biome can be result of an imbalance in sampling effort among the different phytophysognomies of the Serra do Ouro Branco, since the higher sampling effort was conducted with pitfalls within forest fragments. Additional sampling in areas of Cerrado phytophysognomy, like *Campos Rupestres*, may further increase the species richness of the region.

Some species such as *H. mabouia*, *T. torquatus*, *T. merianae* and *A. ameiva*, seem to benefit from human intervention, especially the opening of forest clearings. However, species of the families Gymnophthalmidae, Leiosauridae and Polychrotidae, or 47% of the species in this study, are considered strictly forest species (Jackson 1978; Eisemberg et al. 2004; Verdade and Dixo 2006). Any problems associated with changes in vegetation can cause changes to or even extinction of populations of these lizards which are sensitive to microclimatic changes (Román-Cuesta and Martínez-Vivalta 2006).

No lizard from Serra do Ouro Branco is included in the list of endangered species from the State of Minas Gerais (COPAM 2010), Brazil (MMA 2003) or in the IUCN (2014). The species *Urostrophus vautieri* and *Enyalius perditus* are considered, respectively, vulnerable and possibly threatened, in the list of species of threatened fauna for the state of Rio Grande do Sul (Marques et al. 2002) and for the state of Rio de Janeiro (Rocha et al. 2000). Even though these species are not considered endangered, given the rarity of existing studies for this group, it is of utmost importance to preserve their communities in the various existing habitats (Rodrigues 2005).

Key to the lizards from Serra do Ouro Branco, municipality of Ouro Branco, South of the Espinhaço Range, Minas Gerais, Brazil

- 1a Forelimbs absent; Vestigial and toe-like hindlimbs ***Ophiodes striatus*** (Figure 2A)
- 1b All limbs present and well developed 2
- 2a Dorsal region of the head covered by small, granular scales; eyelids absent; digits short and dilated ***Hemidactylus mabouia*** (Figure 2B)
- 2b Dorsal region of head with scales of variable sizes, but never granular; eyelids present; digits not dilated 3
- 3a Dorsal region of the head covered by numerous scales, arranged irregularly. 4
- 3b Dorsal region of the head covered relatively few scales arranged in regular plates 9
- 4a Vertebral crest present 5
- 4b Vertebral crest absent 6
- 5a Distinctive enlarged and elongate subocular present; sexual dichromatism absent ***Enyalius bilineatus*** (Figure 2H)
- 5b Enlarged and elongate subocular absent, with orbit bordered below by an arc of small scales; sexual dichromatism present, with adult males uniformly green and females with different shades of brown ***Enyalius perditus*** (Figure 2F and 2G)
- 6a Dermal folds (mite pocket) present on the side of the neck; tail not prehensile; flat triangular scales around anterior part of ear-opening present. 7
- 6b Dermal folds on the side of the neck absent; tail prehensile; flat triangular scales around anterior part of ear-opening absent 8
- 7a One deep and oblique mite pocket on each side of the neck; axillary region without folds; side of the neck with scales similar in size to the adjacent dorsal scales ... ***Tropidurus itambere*** (Figure 3G)
- 7b Two shallow mite pockets on the neck filled with granules; axillary region with two to three folds covered by tiny granules separated by areas with larger scales; side of the neck with scales much smaller than the adjacent dorsal scales ***Tropidurus torquatus*** (Figure 3H)
- 8a Extendable, sac-like gular fan present; cone-shaped eyes with eyelids partially fused; femoral pores present ***Polychrus acutirostris*** (figure 3. B).
- 8b Extendable, sac-like gular fan absent; eyelids not partially fused; femoral pores absent ***Urostrophus vautieri*** (Figure 3A)
- 9a Ventral and dorsal body scales cycloid, similar in shape and size 10
- 9b Ventral scales squarish and distinct from dorsals 11
- 10a One frontoparietal scale ***Notomabuya frenata*** (Figure 3C)
- 10b Two frontoparietal scales ***Aspronema dorsivittatum*** (Figure 3D)
- 11a Nasals separated by frontonasal 12
- 11b Nasals in contact medially 14
- 12a Reduced front limbs, with four digits; ear opening absent ... ***Heterodactylus imbricatus*** (Figure 2E)
- 12b Front limbs with five digits; ear opening present. .. 12

- 13a With 9–10 femoral pores.....
.....***Cercosaura quadrilineatus*** (Figure 2C)
- 13b Femoral pores absent.....
.....***Ecpleopus gaudichaudii*** (Figure 2D)
- 14a Ventral rows: 29–33 transverse and 10 across mid
body; preanal pores absent.....
.....***Ameiva ameiva*** (Figure 3E)
- 14b Dorsum with black transverse bands; Ventral
rows: 30–38 transverse and 133–172 scales across
midbody; preanal pores present.....
.....***Tupinambis merianae*** (Figure 3F)

ACKNOWLEDGEMENTS: We thank the Fundação de Amparo a Pesquisa de Minas Gerais (FAPEMIG) for financial support and for the undergraduate grants for the project. We thank the Federal University of Ouro Preto for financial support for the development of this work, the translation of this paper, and for the transport to Ouro Branco. We thank the residents of the municipality of Ouro Branco, especially the residents of the community of Itatiaia, for their support and cooperation during the execution of the project. We thank Leticia, Mariana, Caryne Braga, Lorena Pinto, Valeria Braga, Rafael, Lívia, Ana Clara, Ceres and all the staff from LZV who helped in the field collections. We thank Prof. Yasmine Antonini for the help and collaboration. We thank Vinicius São Pedro and Pedro H. Bernardo for critical reviewing the manuscript and Henrique C. Costa for suggestions and providing bibliographic material, and Pedro Navarro for the maps.

LITERATURE CITED

- Almeida-Abreu, P.A. and F.E. Renger. 2002. A Serra do Espinhaço Meridional: um orógeno de colisão do Mesoproterozóico. *Revista Brasileira de Geociências* 32(1): 1–14 (http://sbgeo.org.br/pub_sbg/rbg/vol32_down/3201/1139.pdf).
- Assis, V.B. 1999. Introdução às serpentes da “Serra do Cipó” (municípios de Santana do Riacho e Jaboticatubas). *Bios* 7(7): 69–71.
- Ávila-Pires, T.C.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). *Zoologische Verhandelingen* 299: 1–706 (<http://www.repository.naturalis.nl/document/149074>).
- Bell, K.E. and M.A. Donnelly. 2006. Influence of Forest Fragmentation on Community Structure of Frogs and Lizards in Northeastern Costa Rica. *Conservation Biology* 20(6): 1750–1760 (doi: 10.1111/j.1523-1739.2006.00522.x).
- Bérnills, R.S. and H.C. Costa (org.). 2012. *Brazilian Reptiles—List of Species. Version 2012.1*. Available at <http://www.sbherpetologia.org.br/>. Sociedade Brasileira de Herpetologia. Captured on 31 August 2013.
- Bérnills, R.S., C. Nogueira and V.X. Silva. 2009. Biota minas—diagnóstico do conhecimento de vertebrados: Répteis; pp. 252–278, in: G.M. Drummond, C.S. Martins, M.S. Greco and F. Vieira (ed.). *Biota minas—diagnóstico do conhecimento sobre a biodiversidade no estado de Minas Gerais*. Belo Horizonte, MG: Fundação Biodiversitas (<http://www.biodiversitas.org.br/biotaminas/publicacao/biotaminas.pdf>).
- Bertoluci, J., M.A.S. Canelas, C.C. Eisemberg, C.F.S. Palmuti and G.G. Montingelli. 2009. Herpetofauna of Estação Ambiental de Peti, an Atlantic Rainforest fragment of Minas Gerais State, southeastern Brazil. *Biota Neotropica* 9(1): 147–175 (doi: 10.1590/S1676-06032009000100017).
- Borges-Martins, M. (1998): *Revisão taxonômica e sistemática filogenética do gênero Ophiodes Wagler, 1828 (Sauria, Anguillidae, Diploglossinae)*. PhD Thesis, Porto Alegre, Pontifícia Universidade Católica do Rio Grande do Sul. 239 pp.
- Carranza, S. and E.N. Arnold. 2006. Systematics, biogeography, and evolution of *Hemidactylus* geckos (Reptilia: Gekkonidae) elucidated using mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 38(2006): 531–545 (doi: 10.1016/j.ympev.2005.07.012).
- Carvalho, A.L.G., A.F.B. Araújo and H.R. Silva. 2007. Lagartos da Marambaia, um remanescente insular de Restinga e Floresta Atlântica no Estado do Rio de Janeiro, Brasil. *Biota Neotropica* 7(2): 221–226 (doi: 10.1590/S1676-06032007000200025).
- Cechin, S.Z. and M. Martins. 2000. Eficiência de armadilhas de queda (pitfall traps) em amostragens de anfíbios e répteis no Brasil. *Revista Brasileira de Zoologia* 17(3): 729–740 (doi: 10.1590/S0101-81752000000300017).
- Cicchi, P.J.P., H. Serafim, M.A. Sena, F.C. Centeno and J. Jim. 2009. Atlantic Rainforest herpetofauna of Ilha Anchieta, an island on municipality of Ubatuba, southeastern Brazil. *Biota Neotropica* 9(2): 201–212 (doi: 10.1590/S1676-06032009000200019).
- Colli, G.R., R.P. Bastos and A.F.B. Araújo. 2002. The character and dynamics of the Cerrado herpetofauna; pp. 223–241, in: P.S. Oliveira and R.L. Marquis (eds.). *The Cerrados of Brazil: Ecology and Natural History of a Neotropical Savanna*. New York: Columbia University Press.
- Colwell, R. K. 2013. EstimateS: Statistical Estimation of Species Richness and Shared Species from samples. University of Connecticut, Connecticut.
- Colwell, R. K., A. Chao, N. J. Gotelli, S.-Y. Lin, C. X. Mao, R. L. Chazdon and J. T. Longino. 2011. Models and estimators linking individual-based and sample-based rarefaction, extrapolation, and comparison of assemblages. *Journal of Plant Ecology* 5: 3–21 (doi: 10.1093/jpe/rtr044).
- Condez, T.H., R.J. Sawaya and M. Dixo. 2009. Herpetofauna dos remanescentes de Mata Atlântica da região de Tapiraí e Piedade, SP, sudeste do Brasil. *Biota Neotropica* 9(1): 1–29 (doi: 10.1590/S1676-06032009000100018).
- Conselho de Política Ambiental. COPAM. Deliberação Normativa COPAM nº 147, de 30 de abril de 2010. *Lista das Espécies da Fauna Ameaçadas de Extinção no Estado de Minas Gerais*. Accessible at <http://www.siam.mg.gov.br/sla/download.pdf?idNorma=13192>. Captured on 08 August 2013.
- Costa, H.C., V.D. Fernandes, A.C. Rodrigues and R.N. Feio. 2009. Lizards and amphisbaenians, municipality of Viçosa, State of Minas Gerais, Southeastern Brazil. *Check List* 5(3): 732–745 (<http://www.checklist.org.br/getpdf?SL131-08>).
- Costa, H.C., V.D. Fernandes, D. Vrcibradic and R.N. Feio. 2008. Reptilia, Scincidae, *Mabuya frenata*: Distribution Extension. *Check List* 4(1): 86–88 (<http://www.checklist.org.br/getpdf?NGD016-06>).
- Crump, M.L. and N.J. Scott Jr. 1994. Visual encounter surveys; pp. 84–92, in: W.R. Heyer, M.A. Donnelly, R.W. Mc Diarmid, L.A.C. Hayek and M.S. Foster (ed.). *Measuring and Monitoring Biological Diversity. Standard Methods for Amphibians*. Washington: Smithsonian Institution Press.
- Derby, O.A. 1906. The Serra of Espinhaço. *Journal of Geology* 14(2): 374–401.
- Dias, E.J.R. and C.F.D. Rocha. 2013. *Ecpleopus gaudichaudi* Duméril and Bibron, 1839 (Squamata: Gymnophthalmidae) and *Psychosaura agmosticha* (Rodrigues, 2000) (Squamata: Scincidae): Distribution extension and new records from Atlantic Forest in Bahia state, Brazil. *Check List* 9(3): 607–609 (<http://www.checklist.org.br/getpdf?NGD173-11>).
- Dixo, M. and J.P. Metzger. 2009. Are corridors fragment size and forest structure important for leaf-litter lizards in fragmented landscape? *Oryx* 43(3): 435–442 (doi: 10.1017/S0030605309431508).
- Dixo, M. and M. Martins. 2008. Are leaf-litter frogs and lizards affected by edge effects due to forest fragmentation in Brazilian Atlantic forest? *Journal of Tropical Ecology* 24(5): 551–554 (doi: 10.1017/S0266467408005282).
- Dixo, M. and V.K. Verdade. 2006. Herpetofauna de serrapilheira da Reserva Florestal de Morro Grande, Cotia (SP). *Biota Neotropica* 6(2): 1–20 (doi: 10.1590/S1676-06032006000200009).
- Doan, T.M. 2003. A new phylogenetic classification for the gymnophthalmid genera *Cercosaura*, *Pantodactylus* and *Prionodactylus* (Reptilia: Squamata). *Zoological Journal of Linnean Society* 137: 101–115 (doi: 10.1046/j.1096-3642.2003.00043.x). Drummond, G.M., C.S. Martins, A.B.M. Machado, F.A. Sebaio, and Y.I. Antonini. (org.). 2005. *Biodiversidade em Minas Gerais: um Atlas para sua conservação*. 2nd ed. Belo Horizonte: Fundação Biodiversitas. 222 pp.
- Eisemberg, C.C., J. Cassimiro and J. Bertoluci. 2004. Notes on the diet of the rare Gymnophthalmid lizard *Ecpleopus gaudichaudii* from Southeastern Brazil. *Herpetological Review* 35(4): 336–337.
- Estes, R., K. De Queiroz and J. Gauthier. 1988. Phylogenetic relationships within Squamata; pp. 119–281; in: R. Estes and G. Pregill (eds.). *Phylogenetic Relationships of the Lizards Families*. Palo Alto: Stanford University Press.
- Etheridge, R. and E.E. Williams. 1991. A review of the South American lizard genera *Urostrophus* and *Anisolepis* (Squamata: Iguania: Polychridae). *Bulletin of the Museum of Comparative Zoology* 152(5): 317–361 (https://archive.org/details/cbarchive_120876_areviewofthesouthamericanlizar1863).
- Feio, R.N. and U. Caramaschi. 2002. Contribuição ao conhecimento da herpetofauna do nordeste do estado de Minas Gerais, Brasil. *Phyllomedusa* 1(2): 105–111 (<http://www.phyllomedusa.esalq.usp.br/articles/volume1/number2/12105111.pdf>).
- Forlani, M.C., P.H. Bernardo, C.B.F. Haddad and H. Zaher. 2010. Herpetofauna of the Carlos Botelho State Park, São Paulo State, Brazil. *Biota Neotropica* 10(3): 265–309 (doi:10.1590/S1676-06032010000300028).
- Frost, D.R., R. Etheridge, D. Janies and T.A. Titus. 2001. Total evidence, sequence alignment, evolution of polychrotid lizards, and a reclassification of the *Iguania* (Squamata, Iguania). *American Museum of Novitates* 3343: 1–38 (doi: 10.1206/0003-0082(2001)343<0001:TE SAE0>2.0.CO;2).
- Gallardo, J.M. 1966. Las especies argentinas del género *Ophiodes* Wagler (Anguillidae, Sauria). *Revista Museo Argentino Ciencias Naturales Bernardino Rivadavia* 9(6): 123–146.
- Gamble, T., A.M. Bauer, E. Greenbaum and T.R. Jackman. 2008. Out

- of the blue: a novel, trans-Atlantic clade of geckos (Gekkota, Squamata). *Zoologica Scripta* 37(4): 355–366 (doi: 10.1111/j.1463-6409.2008.00330.x).
- Gardner, T.A., J.Barlow and C.A. Peres. 2007. Paradox, presumption and pitfalls in conservation biology: The importance of habitat change for amphibians and reptiles. *Biological Conservation* 138: 166–179 (doi: 10.1016/j.biocon.2007.04.017).
- Gasparini, J.L., D.A. Koski and P.L.V. Peloso. 2010. Reptilia, Squamata, Leiosauridae, *Urostrophus vautieri* Duméril and Bribon, 1837: Distribution extension, new state record, and geographic distribution map. *Check List* 6(3): 432–433 (<http://www.checklist.org.br/getpdf?NGD113-09>)
- Gibbons, J.W., D.E. Scott, T.J. Ryan., K.A. Buhmann, T.D. Tuberville, B.S. Metts, J.L. Greene, T. Mills, Y. Leiden, S. Poppy and C.T. Winne. 2000. The global decline of reptiles, deja-vu amphibians. *Bioscience* 50(8): 653–667. (doi: 10.1641/0006-3568(2000)050[0653:TGDORD]2.0.CO;2)
- Giulietti, A.M., J.R. Pirani and R.M. Harley. 1997. Espinhaço range region; pp. 397–404, in: S.D. Davis (ed). *Centres of Plant Diversity: A Guide and Strategy for Their Conservation*. Cambridge, Reino Unido: WWF/IUCN.
- Hedges, S.B. and C.E. Conn. 2012. A new skink fauna from Caribbean islands (Squamata, Mabuyidae, Mabuyinae). *Zootaxa* 3288: 1–244. (<http://www.mapress.com/zootaxa/2012/f/z03288p244f.pdf>)
- Huey, R.B., E.R. Pianka and T.W. Schoener. 1983. Introduction; 1–6 pp, in: R.B. Huey, E.R. Pianka and T.W. Schoener (eds.). *Lizard Ecology: Studies of a Model Organism*. Cambridge: Harvard University Press.
- IUCN 2014. *IUCN Red List of Threatened Species. Version 2014.2*. Electronic database accessible at <http://www.iucnredlist.org/>. Captured on 5 August 2014.
- Jackson, J.F. 1978. Differentiation in the genera *Enyalius* and *Strobilurus* (Iguanidae): implications for Pleistocene climatic changes in eastern Brazil. *Arquivos de Zoologia* 30(1): 1–79 (doi:10.11606/issn.2176-7793.v30i1p1-79).
- Jellinek, S., D.A. Driscoll and J.B. Kirkpatrick. 2004. Environmental and vegetation variables have a greater influence than habitat fragmentation in structuring lizard communities in remnant urban bushland. *Austral Ecology* 29(3): 294–304 (doi:10.1111/j.1442-9993.2004.01366.x).
- Juncá, F.A. 2005. Anfíbios e Répteis; pp. 339–376, in: F.A. Juncá, L. Funch and W. Rocha (ed.). *Biodiversidade e Conservação da Chapada Diamantina*—Série Biodiversidade. Brasília: Ministério do Meio Ambiente.
- Kawashita-Ribeiro, R.A. and R.W. Ávila. 2008. Reptilia, Squamata, *Polychrus* spp.: New record, range extensions, and distribution map in the state of Mato Grosso, Brazil. *Check List* 4(3): 362–365 (<http://www.checklist.org.br/getpdf?NGD308-08>).
- Kitchener, D.J., A. Chapman., J. Dell, B.G. Muir and M. Palmer. 1980. Lizard assemblages and reserve size and structure in the Western Australian wheatbelt - some implications for conservation. *Biological Conservation* 17(1): 25–62 (doi:10.1016/0006-3207(80)90024-5).
- Lehtinen, R.M., J-B Ramanamanjato and J.G. Raveloarison. 2003. Edge effects and extinction proneness in a herpetofauna from Madagascar. *Biodiversity and Conservation* 12: 1357–1370 (doi:10.1023%2FA%3A1023673301850).
- Linares, A. M., and P. C. Eterovick. 2013. Herpetofaunal Surveys support successful reconciliation ecology in secondary and human-modified habitats at the Inhotim Institute, Southeastern Brasil. *Herpetologica* 69 (2): 237–256 (doi:10.1655/HERPETOLOGICA-D-12-00030).
- Marques, A.A.B., C.S. Fontana, E. Vélez, G.A. Bencke, M. Schneider and R.S. Reis. 2002. *Lista de Referência da Fauna Ameaçada de Extinção no Rio Grande do Sul*. Decreto no. 41.672, 11 de junho de 2002. Porto Alegre: FZB/MCT-PUCRS/PANGEA. 52 pp. (http://www.fzb.rs.gov.br/upload/1396360907_fauna_ameacada.pdf).
- Ministério do Meio Ambiente. 2003. Instrução Normativa nº 003, de 26 de Maio de 2003.
- Mittermeier, R.A., P.R. Gil, M. Hoffmann, J. Pilgrim, T. Brooks, C.G. Mittermeier, J. Lamourux and G.A.B. Fonseca. 2004. *Hotspots Revisited: earth's Biologically Richest and Most Endangered Terrestrial Ecoregions*. P. 390. Mexico: CEMEX. (<http://www.conservation.org.br/publicacoes/files/HotspotsRevisitados.pdf>).
- Moura, M.R., A.P. Motta, V.D. Fernandes and R.N. Feio. 2012. Herpetofauna da Serra do Brigadeiro, um remanescente da Mata Atlântica, em Minas Gerais, Sudeste do Brasil. *Biota Neotropica* 12(1): 209–235 (doi:10.1590/S1676-06032012000100017).
- Nogueira, C. C. 2006. Diversidade e padrões de distribuição da fauna de lagartos do Cerrado. Tese de Doutorado. USP-Instituto de Biociências São Paulo.
- Novelli, I.A., P.S. Lucas and R.C. Santos. 2011. Reptila, Squamata, Gymnophthalmidae, *Heterodactylus imbricatus* Spix, 1825: Filling gaps in the state of Minas Gerais. *Check List* 7(1): 30–31 (<http://www.checklist.org.br/getpdf?NGD017-10>).
- Novelli, I.A., P.S. Lucas., R.G. Carvalho., R.C. Santos and B.M. Sousa. 2012. Lizards of the Cerrado areas in the Reserva Biológica Unilavras-Boqueirão, Ingaí, Minas Gerais Southern, Brazil. *Biota Neotropica* 12(3): 147–153 (doi:10.1590/S1676-06032012000300017).
- Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, R. B. O'hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, and H. Wagner. 2010. Vegan: community ecology package. R package version 1.17-3 (<http://cran.r-project.org/web/packages/vegan/index.html>).
- Peters, J.A. and R. Donoso-Barros. 1970. Catalogue of the Neotropical Squamata. Part II. Lizards and Amphisbaenians. *Bulletin of the United States National Museum* 297(2): 1–293.
- Pianka, E. R. 1973. The structure of lizards communities. *Annual review of ecology and systematics* 4: 53–74 (doi:10.1146/annurev.es.04.110173.000413).
- Pyron, R.A., F.T. Burbrink and J.J. Wiens. 2013. A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evolutionary Biology* 13: 93 (doi:10.1186/1471-2148-13-93).
- Quintela, F.M., D. Loebmann and N.M. Gianuca. 2006. Répteis continentais do município de Rio Grande, Rio Grande do Sul, Brasil. *Biociências* 14(2): 180–188 (<http://revistaseletronicas.pucrs.br/fo/ojs/index.php/fabio/article/view/237>).
- R Development Core Team. 2013. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Viena. (<http://www.gbif.org/resources/2585>).
- Ramos, L.O. and A.L. Silveira. 2011. New Record and distribution mape of *Ecleopus gaudichaudii* (Reptilia, Squamata, Gymnophthalmidae) in southeastern Brazil. *Herpetology Notes* 4: 173–176 (http://www.herpetologynotes.seh-herpetology.org/Volume4_PDFs/Ramos_Silveira_Herpetology_Notes_Volume4_pages173-176.pdf).
- Recoder, R. and C. Nogueira. 2007. Composição e diversidade de répteis na região sul do Parque Nacional Grande Sertão Veredas, Brasil Central. *Biota Neotropica* 7(3): 267–278. (doi:10.1590/s1676-06032007000300029).
- Reinhardt, J. and Lutken, C. 1861. Bidrag til kundskab om Brasiliens Padder og krybdyr Viedenskabelige Meddelesler fra den Naturhistorisk. *Forening Kjöbenhavn* 10–15: 143–242, 6 pls.
- Rocha, C.F.D., D. Vrcibradic, J.J. Vicente and M. Cunha-Barros. 2003. Helminths infecting *Mabuya dorsivittata* (Lacertilia, Scincidae) from a high-altitude habitat in Itatiaia National Park, Rio de Janeiro State, Southeastern Brazil. *Brazilian Journal of Biology* 63(1): 129–132. (doi:10.1590/s1519-69842003000100017).
- Rocha, C.F.D., M. Van-Sluys, G. Puorto, R. Fernandes, J.D. Barros-Filho, R.R.E. Silva, F.A. Neo and E.A. Melgarejo. 2000. Répteis; pp. 79–87, in: H.G. Bergallo, C.F.D. Rocha, M.A.S. Alves and M. Van-Sluys (ed.). *A Fauna Ameaçada de Extinção do Estado do Rio de Janeiro*. Rio de Janeiro: Editora UERJ.
- Rocha, C.F.D., L.A. Anjos and H.G. Bergallo. 2011. Conquering Brazil: the invasion by the exotic gekkonid lizard *Hemidactylus mabouia* (Squamata) in Brazilian natural environments. *Zoologia* 28(6): 747–754 (doi:10.1590/S1984-46702011000600007).
- Rödger, D., M. Solé and W. Böhme. 2008. Predicting the potential distributions of two alien invasive Housegeckos (Gekkonidae: *Hemidactylus frenatus*, *Hemidactylus mabouia*). *North-western Journal of Zoology* 4(2): 236–246.
- Rodrigues, M.T. 1987. Sistemática, ecologia e zoogeografia dos Tropicurus do grupo *torquatus* ao sul do Rio Amazonas (Sauria, Iguanidae). *Arquivos de Zoologia* 31(3): 105–230 (doi:10.11606/issn.2176-7793.v31i3p105-230).
- Rodrigues, M.T. 2005. Conservação dos répteis brasileiros: os desafios para um país megadiverso. *Megadiversidade* 1(1): 87–94 (http://www.conservation.org.br/publicacoes/megadiversidade/13_Rodrigues.pdf).
- Rodrigues, M.T., K.C.M. Pellegrino, M. Dixo, V.K. Verdade, D. Pavan, A.J.S. Argolo and J.W. Sites Jr. 2007. A new genus of microteiid lizard from the Atlantic Forests of State Bahia, Brazil, with a new generic name for *Colobosaura mentalis mentalis*, and a discussion of relationships among the *Heterodactylini* (Squamata, Gymnophthalmidae). *American Museum Novitates* 3565: 1–27 (<http://hdl.handle.net/2246/5856>).
- Rodrigues, M.T., M.A. Freitas and T.F.S. Silva. 2009. New Species of earless lizard genus *Heterodactylus* (Squamata: Gymnophthalmidae) from the highlands of Chapada Diamantina, State of Bahia, Brazil. *Journal of Herpetology* 43(4): 605–611 (doi:10.1670/08-133.1).
- Román-Cuesta R. M and J. Martínez-Vilalta. 2006) Effectiveness of protected areas in mitigating fire within their boundaries: case study of Chiapas, Mexico. *Conservation Biology* 20:1074–1086 (doi: 10.1111/j.1523-1739.2006.00478.x).
- Santos, R.C., P.S. Lucas., B.M. Sousa and I.A. Novelli. 2009. Reptilia, Squamata, Leiosauridae, *Urostrophus vautieri*: Distribution extension and geographic distribution map. *Check List* 5(3): 533–536 (<http://www.checklist.org.br/getpdf?NGD042-09>).
- Sartorius, S.S., L.J. Vitt and G.R. Colli. 1999. Use of naturally and

- anthropogenically disturbed habitats in Amazonian rainforests by the teiid lizard *Ameiva ameiva*. *Biological Conservation* 90(2): 91–101 (doi:10.1016/S0006-3207(99)00019-1).
- Sazima, I. and C.F.B. Haddad. 1992. Répteis da Serra do Japi: notas sobre História Natural; pp. 212–236, in: L.P.C. Morellato (ed.). *História Natural da Serra do Japi: ecologia e preservação de uma área florestal do sudeste do Brasil*. Campinas: UNICAMP/FAPESP.
- São Pedro and M.R.S. Pires. 2009. As Serpentes da região de Ouro Branco, Extremo Sul da Cadeia do Espinhaço, Minas Gerais. *Ceres* 56 (2): 166–171 (<http://www.ceres.ufv.br/ceres/revistas/V56N002P50809.pdf>).
- Silveira, A.L., M.R.S. Pires and G.A. Cotta. 2010. Serpentes de uma área de transição entre o Cerrado e a Mata Atlântica, no sudeste do Brasil. *Arquivos do Museu Nacional* 68(1-2): 79–110.
- Sousa, B.M., A.E.R. Nascimento, S.C. Gomides, C.H. Varela Rios, A.H. Hudson and I.A. Novelli. 2010. Reptiles in fragments of Cerrado and Atlantic Forest at the Campo das Vertentes, MinasGerais State, Southeastern Brazil. *Biota Neotropica* 10(2): 129–138 (doi:10.1590/S1676-06032010000200016).
- Sousa, B.M., S.C. Gomides, A.A. Hudson, L.B. Ribeiro and I.A. Novelli. 2012. Reptiles of the municipality of Juiz de Fora, Minas Gerais state, Brazil. *Biota Neotropica* 12(3): 35–49 (doi:10.1590/S1676-06032012000300002).
- Uetz, P and J. Hosek (eds.). 2013. *The Reptile Database*. Accessible at <http://www.reptile-database.org>. Captured on 09 July 2013 (<http://www.reptile-database.org/>).
- Vanzolini, P.E. 1948. Sobre um novo Pantadactylus do Estado de Minas Gerais (Sauria, Teeidae). *Papeis avulsos do Departamento de Zoologia* 7 (27): 337–340.
- Vanzolini, P.E. 1978. On South American Hemidactylus (Saurua, Gekkonidae). *Papeis Avulsos de Zoologia* 31 (20): 307–343.
- Vanzolini, P.E. 1983. Guiano-Brasilian *Polychrus*: distribution and speciation (Sauria: Iguanidae); pp. 118–131, in: A.G.J. Rhodin and K.Miyata (ed.). *Advances in Herpetology and Evolutionary Biology*. Cambridge: Museum of Comparative Zoology.
- Vanzolini, P.E. 1986. Addenda and corrigenda to the Catalogue of Neotropical Squamata. *Smithsonian Herpetological Information Service* 70: 1-25 (<https://repository.si.edu/handle/10088/8291>).
- Vanzolini, P.E., A.M Ramos-Costa and L.J. Vitt. 1980. *Répteis das Caatingas*. Rio de Janeiro: Academia Brasileira de Ciências. 161 pp.
- Veloso, H.P., A.L.R. Rangel-Filho and J.C.A. Lima. 1991. Classificação da vegetação brasileira adaptada a um sistema universal. Rio de Janeiro: *Instituto Brasileiro de Geografia e Estatística* (IBGE). (<http://biblioteca.ibge.gov.br/visualizacao/monografias/GEBIS%20-%20RJ/classificacaovegetal.pdf>).
- Vitt, L.J. and G.R. Colli. 1994. Geographical ecology of a Neotropical lizard: *Ameiva ameiva* (Teiidae) in Brazil. *Canadian Journal of Zoology* 72(11): 1986–2008 (doi:10.1139/z94-271).
- Vitta, F.A. 2002. Diversidade e conservação da flora nos campos rupestres na Cadeia do Espinhaço em Minas Gerais; pp. 90–94, in: L.A. Araujo (ed.). *Biodiversidade, Conservação e Uso Sustentável da Flora do Brasil*. Sociedade Botânica do Brasil, Recife: Imprensa Universitária.

RECEIVED: April 2014

ACCEPTED: August 2014

PUBLISHED ONLINE: December 2014

EDITORIAL RESPONSIBILITY: Ross MacCulloch

APPENDIX 1: Voucher specimens from Serra do Ouro Branco, state of Minas Gerais, Brazil

Ameiva ameiva: LZV 494S, 616S. *Aspronema dorsivittatum*: MZUFV-530. *Cercosaura quadrilineatus*: LZV 003S-L. *Ecpleopus gaudichaudii*: LZV 973S, 991S, 992S, 1000S, 1006S, 1007S, 1008S, 1009S, 1010S, 1024S, 1041S, 1044S, 1049S, 1050S, 1051S, 1063S, 1064S, 1068S, 1069S, 1090S, 1091S, 1101S, 1111S, 1112S, 1113S, 1114S, 1115S, 1116S, 1148S. *Enyalius bilineatus*: LZV 943S, 971S, 1060S, 1065S. *Enyalius perditus*: LZV 939S, 940S, 942S, 944S, 966S, 967S, 968S, 976S, 977S, 978S, 993S, 995S, 996S, 997S, 1005S, 1038S, 1039S, 1040S, 1042S, 1043S, 1045S, 1047S, 1048S, 1057S, 1059S, 1062S, 1067S, 1070S, 1071S, 1075S, 1076S, 1077S, 1079S, 1080S, 1081S, 1082S, 1083S, 1085S, 1086S, 1087S, 1098S, 1099S, 1117S, 1118S, 1109S, 1110S. *Hemidactylus mabouia*: LZV 207S. *Heterodactylus imbricatus*: LZV 941S, 974S, 990S, 999S, 1017S, 1019S, 1052S, 1058S, 1119S 1100S. *Notomabuya frenata*: LZV 1061S, 1066S, 002S-L. *Ophiodes striatus*: LZV 746S, 811S, 937S, 938S. *Polychrus acutirostris*: LZV 028S-L. *Tropidurus itambere*: LZV 873S. *Tropidurus torquatus*: LZV 232S. *Tupinambis merianae*: LZV 994S. *Urostrophus vautieri*: LZV 1046S.

APPENDIX 2: *Ophiodes striatus*: Peters and Donoso 1970, Galardo 1966; *Hemidactylus mabouia*: Vanzolini et al. 1980, Avila-Pires 1995, Carranza and Arnold 2006, *Cercosaura quadrilineatus*: Vanzolini 1948, Doan 2003; *Ecpleopus gaudichaudii*: Ramos and Silveira 2011, Dias and Rocha 2013; *Heterodactylus imbricatus*: Dixo and Verdade 2006, Rodrigues et al. 2007, 2009, Novelli et al. 2011; *Enyalius bilineatus*: Jackson 1978; *Enyalius perditus*: Jackson 1978; *Urostrophus vautieri*: Etheridge and Williams 1991, Sazima and Haddad 1992, Santos et al. 2009, Gasparini et al. 2010; *Polychrus acutirostris*: Vanzolini et al. 1980, Vanzolini 1983, Kawashita-Ribeiro and Avila 2008; *Aspronema dorsivittatum*: Recorder and Nogueira 2007, Quintela et al. 2006, Rocha et al. 2003; *Notomabuya frenata*: Peters and Danoso-Barros 1970, Recorder and Nogueira 2007, Costa et al. 2008, Sousa et al. 2010; *Ameiva ameiva*: Vanzolini 1986, Vitt and Colli 1994, Sartorius et al. 1999; *Tupinambis merianae*: Sazima and Haddad 1992, Vanzolini et al. 1980, Ávila-Pires 1995; *Tropidurus torquatus*: Rodrigues 1987; *Tropidurus itambere*: Rodrigues 1987.